

Probing High-Energy Physics with High-precision QED measurements ¹

Stephan Narison

Laboratoire de Physique Mathématique
Université de Montpellier II
Place Eugène Bataillon
34095 - Montpellier Cedex 05, France
Email: qcd@lpm.univ-montp2.fr

Abstract

I summarize our *self-contained* determinations of the lowest order hadronic contributions [1, 2] to the anomalous magnetic moments $a_{\mu,\tau}$ of the muon and tau leptons, the running QED coupling $\alpha(M_Z)$ and the muonium hyperfine splitting ν . Using an average estimate of the light-by light scattering contribution: $a_\mu(LL) = 85(18) \times 10^{-11}$, we deduce: $a_\mu^{SM} = 116\,591\,861(78) \times 10^{-11}$, $a_\tau^{SM} = 117\,759(7) \times 10^{-8}$, giving: $a_\mu^{SM} - a_\mu^{exp} = 162(170) \times 10^{-11}$. We also obtain: $\alpha^{-1}(M_Z) = 128.926(25)$ and the Fermi energy splitting: $\nu_F^{SM} = 4\,459\,031\,783(229)$ Hz. Lower bounds on some new physics are given, while ν_F^{SM} leads e.g. to $m_\mu/m_e = 206.768\,276(11)$ in remarkable agreement with the data.

1 Introduction

QED is at present the gauge theory where perturbative calculations are the most precise known today. Therefore, accurate measurements of QED processes are expected to give strong constraints on different electroweak models and to reveal some eventual deviations from the standard model (SM) predictions ². In the following, I will discuss the effects of the hadronic and QCD contributions to three classical QED processes which are: the anomalous magnetic moment of the muon and tau leptons, the running QED coupling $\alpha(M_Z)$ and the muonium hyperfine splitting ν . These hadronic contributions are one of the main sources of uncertainties into these processes. Using a dispersion relation, it is remarkable to notice that the different lowest order hadronic contributions for these three processes can be expressed in a closed form as a convolution of the $e^+e^- \rightarrow$ hadrons cross-section $\sigma_H(t)$ with a QED kernel function $K(t)$ which depends on each observable:

$$\mathcal{O}_{\text{had}} = \frac{1}{4\pi^3} \int_{4m_\pi^2}^{\infty} dt K_{\mathcal{O}}(t) \sigma_H(t) , \quad (1)$$

where:

$$\mathcal{O}_{\text{had}} \equiv a_{l,\text{had}} , \quad \Delta\alpha_{\text{had}} \times 10^5 \quad \text{or} \quad \Delta\nu_{\text{had}} . \quad (2)$$

¹Talk presented at the 1st High-Energy Physics Madagascar International Conference Series (HEP-MAD'01), 27th sept-5th Oct. 2001, Antananarivo (to be published by World Scientific, Singapore).

²For general discussions on astroparticle physics and supersymmetric models, see e.g. [3, 4].

- For the anomalous magnetic moment $a_{l,\text{had}}$, $K_{a_l}(t \geq 0)$ is the well-known kernel function [5]:

$$K_{a_l}(t) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (t/m_l^2)(1-x)} , \quad (3)$$

where m_l is the lepton mass. It behaves for large t as:

$$K_{a_l}(t \gg m_l^2) \simeq \frac{m_l^2}{3t} . \quad (4)$$

- For the QED running coupling $\Delta\alpha_{\text{had}} \times 10^5$, the kernel is (see e.g. [6]):

$$K_\alpha(t) = \left(\frac{\pi}{\alpha}\right) \left(\frac{M_Z^2}{M_Z^2 - t}\right) , \quad (5)$$

where $\alpha^{-1}(0) = 137.036$ and $M_Z = 91.3$ GeV. It behaves for large t like a constant.

- For the muonium hyperfine splitting $\Delta\nu_{\text{had}}$, the kernel function is (see e.g [7]):

$$K_\nu = -\rho_\nu \left[(x_\mu + 2) \ln \frac{1+v_\mu}{1-v_\mu} - \left(x_\mu + \frac{3}{2}\right) \ln x_\mu \right] \quad (6)$$

where:

$$\rho_\nu = 2\nu_F \frac{m_e}{m_u} , \quad x_\mu = \frac{t}{4m_\mu^2} \quad v_\mu = \sqrt{1 - \frac{1}{x_\mu}} , \quad (7)$$

and we take (for the moment) for a closed comparison with [7]³, the value of the Fermi energy splitting:

$$\nu_F = 445\,903\,192\,0.(511)(34) \text{ Hz} . \quad (8)$$

It behaves for large t as:

$$K_\nu(t \gg m_\mu^2) \simeq \rho_\nu \left(\frac{m_\mu^2}{t}\right) \left(\frac{9}{2} \ln \frac{t}{m_\mu^2} + \frac{15}{4}\right) . \quad (9)$$

The different asymptotic behaviours of these kernel functions will influence on the relative weights of different regions contributions in the evaluation of the above integrals.

2 Input and Numerical Strategy

The different data input and QCD parametrizations of the cross-section $\sigma_H(t)$ have been discussed in details in [1] (herereferred as SN1) and corresponding discussions will not be repeated here. The sources of these data are quoted in the last column of Table 1 from SN1 and Table 2 from [2] (herereferred as SN2) are classified according to the estimate in different regions. We shall only sketched briefly the numerical strategy here:

- Our result from the $I = 1$ isovector channel below 3 GeV^2 is the mean value of the one using τ -decay and e^+e^- data. In both cases, we have used standard trapezoidal rules and/or least square fits of the data in order to avoid theoretical model dependence parametrization of the pion form factor. Correlations among different data have been taken in the compilations of [8] used in this paper. In the region $(0.6 - 0.8) \text{ GeV}^2$ around the ω - ρ mixing, we use in both cases e^+e^- data in order to take properly the $SU(2)_F$ mixing. The $SU(2)$ breaking in the remaining regions are taken into account by making the average of the two results from τ -decay and e^+e^- and by adding into the errors the distance between this mean central value with the one from each data.

³In the next section, we shall extract this value from the analysis.

- For the $I = 0$ isoscalar channel below 3 GeV^2 , we use the contributions of the resonances $\omega(782)$ and $\phi(1020)$ using narrow width approximation (NWA). We add to these contributions, the sum of the exclusive channels from 0.66 to 1.93 GeV^2 . Above 1.93 GeV^2 , we include the contributions of the $\omega(1.42)$, $\omega(1.65)$ and $\phi(1.68)$ using a Breit-Wigner form of the resonances.
- For the heavy quarkonia, we include the contributions of known J/ψ (1S to 4.415) and Υ (1S to 11.02) families and use a NWA. We have added the effect of the $t\bar{t}$ bound state using the leptonic width of $(12.5 \pm 1.5) \text{ keV}$ given in [9].
- Away from thresholds, we use perturbative QCD plus negligible quark and gluon condensate contributions, which is expected to give a good parametrization of the cross-section. These different expressions are given in SN1. However, as the relative rôle of the QCD continuum is important in the estimate of $\Delta\alpha_{\text{had}}$, we have added, to the usual Schwinger interpolating factor at order α_s for describing the heavy quark spectral function, the known $\alpha_s^2 m_Q^2/t$ corrections given in SN1. However, in the region we are working, these corrections are tiny.
- On the Z -mass, the integral for $\Delta\alpha_{\text{had}}$ has a pole, such that this contribution has been separated in this case from the QCD continuum. Its value comes from the Cauchy principal value of the integral.

3 Lowest order hadronic contributions

3.1 Muon and tau anomalies

We show in Table 1 the details of the different hadronic contributions from each channels and from different energy regions for the muon and tau anomalies. Taking the average of the results in Table 1 and adding further systematics due to an eventual deviation from the CVC assumption and from the choice of the QCD continuum threshold for the light flavours, one deduce the final estimate from an average of the e^+e^- and τ -decay data [1]:

$$a_\mu^{\text{had}}(l.o) = 7020.6(75.6) \times 10^{-11}, \quad a_\tau^{\text{had}}(l.o) = 353.6(4.0) \times 10^{-8}, \quad (10)$$

- The main error (80% when added quadratically) in our previous determinations comes from the ρ -meson region below 0.8 GeV^2 . Hopefully, improved measurements of this region are feasible in the near future.
- The second source of errors comes from the region around M_τ for the inclusive τ -decay and between 1 GeV to M_τ for the e^+e^- data. These errors are about half of the one from the region below 0.8 GeV^2 in most different determinations. They can be reduced by improved measurements of inclusive τ -decay near M_τ ($I = 1$) and by improving the measurements of the odd multi-pions and $\bar{K}K$, $\bar{K}K\pi, \dots$ channels in the $I = 0$ channels from e^+e^- data.
- The contributions of the whole region above 3 GeV^2 induce much smaller errors (7% of the total). There is a quite good consensus between different determinations in this energy region.
- These predictions agree within the errors with previous predictions quoted in SN1 [1] and recent estimates given in [6]–[11].

3.2 Running QED coupling

Using the same data as for the anomalous magnetic moment, one can deduce from Table 2:

$$\Delta\alpha_{\text{had}} = 2763.4(16.5) \times 10^{-5}. \quad (11)$$

Also a detailed comparison of each region of energy with the most recent work of [9] shows the same features (agreement and slight difference) like in the case of a_μ in SN1 due to the slight difference

in the parametrization of the data and spectral function. However, the final results are comparable. Finally, one can remark that due to the high-energy constant behaviour of the QED kernel function in this case, the low-energy region is no longer dominating. For a_μ , the contribution of the ρ -meson below 1 GeV is 68% of the total contribution, while the sum of the QCD continuum is only 7.4% (see e.g. SN1). Here the situation is almost reversed: the contribution of the ρ -meson below 1 GeV is only 2%, while the sum of the QCD-continuum is 73.6%.

Table 1: Determinations of $a_l^{\text{had}}(l.o)$ using combined e^+e^- and inclusive τ decay data (2nd and 4th columns) and averaged e^+e^- data (3rd column).

Region in GeV ²	$\mathbf{a}_\mu^{\text{had}}(\text{l.o}) \times 10^{11}$	$\mathbf{a}_\tau^{\text{had}}(\text{l.o}) \times 10^8$	Data input
	$\tau + \mathbf{e}^+ \mathbf{e}^-$	$\mathbf{e}^+ \mathbf{e}^-$	$\tau + \mathbf{e}^+ \mathbf{e}^-$
Light Isovector			
$4m_\pi^2 \rightarrow 0.8$	4794.6 ± 60.7	4730.2 ± 99.9	165.8 ± 1.5 [8, 12, 13]
$0.8 \rightarrow 2.1$	494.9 ± 15.8	565.0 ± 54.0	28.7 ± 1.1 [12, 13]
$2.1 \rightarrow 3.$	202.0 ± 29.7	175.9 ± 16.0	17.0 ± 2.6 [12, 13]
<i>Total Light I=1</i>	5491.5 ± 69.4	5471.1 ± 114.7	211.5 ± 3.2
Light Isoscalar			
<i>Below 1.93</i>			
ω	387.5 ± 13	387.5 ± 13	15.3 ± 0.5 NWA [14]
ϕ	393.3 ± 9.9	393.3 ± 9.9	21.0 ± 0.5 NWA [14]
$0.66 \rightarrow 1.93$	79.3 ± 18.7	79.3 ± 18.7	4.3 ± 1.1 $\sum$ exclusive [15]
<i>From 1.93 to 3</i>			
$\omega(1.42), \omega(1.65)$	31.3 ± 6.8	31.3 ± 6.8	2.6 ± 0.7 BW [16, 14]
$\phi(1.68)$	42.4 ± 18.2	42.4 ± 18.2	3.8 ± 1.3 BW [16, 17, 14]
<i>Total Light I=0</i>	933.8 ± 31.5	933.8 ± 31.5	47.0 ± 2.0
Heavy Isoscalar			
$J/\psi(1S \rightarrow 4.415)$	87.0 ± 4.7	87.0 ± 4.7	13.08 ± 0.69 NWA [14]
$\Upsilon(1S \rightarrow 11.020)$	0.95 ± 0.04	0.95 ± 0.04	0.23 ± 0.01 NWA [14]
<i>Total Heavy I=0</i>	88.0 ± 4.7	88.0 ± 4.7	13.3 ± 0.7
QCD continuum			
$3. \rightarrow (4.57)^2$	407.0 ± 2.3	407.0 ± 2.3	49.4 ± 0.3 (u, d, s)
$(4.57)^2 \rightarrow (11.27)^2$	95.3 ± 0.5	95.3 ± 0.5	27.3 ± 0.1 (u, d, s, c)
$(11.27)^2 \rightarrow 4M_t^2$	20.5 ± 0.1	20.5 ± 0.1	5.87 ± 0.01 (u, d, s, c, b)
$4M_t^2 \rightarrow \infty$	$\approx 0.$	$\approx 0.$	$\approx 0.$ (u, d, s, c, b, t)
<i>Total QCD Cont.</i>	522.8 ± 2.4	522.8 ± 2.4	82.6 ± 0.3
	7036.1(76.4)	7015.7(119.1)	354.4(3.8)

For this reason, improvement due to the new Novosibirsk e^+e^- data [18] in the low-energy region is not significant, as we have explicitly checked. At present, new BES data [19] in the J/ψ region are also available, which can be alternatively used. Below the J/ψ resonances, the BES data are in excellent agreement with the QCD parametrization to order α_s^3 used here for 3 flavours, justifying the accuracy of your input. Above the J/ψ resonances, the parametrization used here (sum of narrow resonances +QCD continuum away from thresholds) can also be compared with these data. One can notice that, in the resonance regions, the BES data are more accurate than previous ones, which may indicate that our quoted errors in Table 1 for the J/ψ family contributions are overestimated. In addition, the threshold of the QCD continuum which we have taken above the J/ψ resonances, matches quite well with the one indicated by the BES data. Our estimate of $\Delta\alpha_{\text{had}}$ is compared with recent predictions respectively in Fig. 1 from SN2 [2] (references to the authors are

in [6, 9, 8, 20]) where one can notice a very good agreement.

Table 2: Lowest order determinations of $\Delta\alpha_{\text{had}} \times 10^5$ and $\Delta\nu_{\text{had}}$ [Hz] using combined e^+e^- and inclusive τ decay data (2nd and 4th columns) and averaged e^+e^- data (3rd and 5th columns).

Region in GeV ²	$\Delta\alpha_{\text{had}} \times 10^5$		$\Delta\nu_{\text{had}}$ [Hz]		Data
	$\tau+e^+e^-$	e^+e^-	$\tau+e^+e^-$	e^+e^-	
Light Isovector					
$4m_\pi^2 \rightarrow 0.8$	314.5 ± 2.3	302.7 ± 7.1	152.9 ± 1.8	148.4 ± 3.1	[8, 12, 13]
$0.8 \rightarrow 2.1$	77.2 ± 3.4	82.0 ± 5.4	12.1 ± 0.5	16.9 ± 1.9	[12, 13]
$2.1 \rightarrow 3.$	62.3 ± 9.2	53.6 ± 4.9	7.8 ± 1.2	6.7 ± 0.6	[12, 13]
<i>Total Light I=1</i>	454.0 ± 10.6	438.2 ± 10.2	172.8 ± 2.2	172.1 ± 3.7	
<i>Average</i>	$446.1 \pm 10.4 \pm 7.9$		$172.5 \pm 3.0 \pm 0.3$		
Light Isoscalar					
<i>Below 1.93</i>					
ω	31.5 ± 1.1		12.7 ± 0.4		NWA [14]
ϕ	52.3 ± 1.2		13.7 ± 0.3		NWA [14]
$0.66 \rightarrow 1.93$	11.6 ± 3.0		2.7 ± 0.7		$\sum$ excl. [15]
<i>From 1.93 to 3</i>					
$\omega(1.42), \omega(1.65)$	9.4 ± 1.4		1.2 ± 0.2		BW [16, 14]
$\phi(1.68)$	14.6 ± 4.6		$1.7 \pm .5$		BW [16, 17, 14]
<i>Total Light I=0</i>	119.0 ± 5.9		32.1 ± 1.0		
Heavy Isoscalar					
$J/\psi(1S \rightarrow 4.415)$	116.3 ± 6.2		4.0 ± 0.2		NWA [14]
$\Upsilon(1S \rightarrow 11.020)$	12.7 ± 0.5		0.1 ± 0.0		NWA [14]
$T(349)$	$-(0.1 \pm 0.0)$		≈ 0		NWA [14, 9]
<i>Total Heavy I=0</i>	128.9 ± 6.2		$4.1 \pm .2$		
QCD continuum					
$3. \rightarrow 4.57^2$	330.1 ± 1.0		$17.5 \pm .1$		(u, d, s)
$4.57^2 \rightarrow 11.27^2$	503.0 ± 1.0		$5.0 \pm .1$		(u, d, s, c)
$11.27^2 \rightarrow (M_Z - 3 \text{ GeV})^2$	2025.7 ± 2.0		1.3 ± 0.0		(u, d, s, c, b)
$(M_Z + 3 \text{ GeV})^2 \rightarrow 4M_t^2$	$-(794.6 \pm 0.6)$		≈ 0		—
Z -pole	$29.2 \pm .5$		≈ 0		ppal value [9]
$4M_t^2 \rightarrow \infty$	$-(24.0 \pm 0.1)$		≈ 0		(u, d, s, c, b, t)
<i>Total QCD Cont.</i>	2069.4 ± 5.2		23.8 ± 1.4		
Final value	2763.4 ± 16.5		232.5 ± 3.2		

3.3 The muonium hyperfine splitting

Our final result for $\Delta\nu_{\text{had}}$ comes from Table 2 of SN2 [2]:

$$\Delta\nu_{\text{had}} = (232.5 \pm 3.2) \text{ Hz} \quad (12)$$

and is shown in Table 3 in comparison with other determinations, where there is an excellent agreement with the most recent determination [7]. Here, due to the $(\ln t)/t$ behaviour of the kernel function, the contribution of the low-energy region is dominant. However, the ρ -meson region contribution below 1 GeV is 47% compared with 68% in the case of a_μ , while the QCD continuum is about 10% compared to 7.4% for a_μ . The accuracy of our result is mainly due to the use of the τ -decay data, explaining the similar accuracy of our final result with the one in [7] using new Novosibirsk data. The agreement with [7] can be understood from the agreement of the averaged correlated e^+e^- and τ -decay data compiled in [8] with the new Novosibirsk data used in [7].




Figure 1: Recent determinations of $\Delta\alpha_{\text{had}}$ and $\alpha^{-1}(M_Z)$. The dashed vertical line is the mean central value.

However, we differ with DH98 [20] in the treatment of the QCD contribution ⁴. For light quarks, QCD is applied in the region where non-perturbative contributions are inessential. For heavy quarks, perturbative QCD is applied far from heavy quark thresholds, where it can be unambiguously used.

Table 3: Recent determinations of $\Delta\nu_{\text{had}}$

Authors	$\Delta\nu_{\text{had}}$ [Hz]
FKM 99 [21]	240 ± 7
CEK 01 [7]	233 ± 3
SN 01 [2]	232.5 ± 3.2

4 Theory of the muon and tau anomalies and new physics

4.1 QED and EW contributions to a_μ

The QED up to 8th order and EW including two-loop corrections ⁵ are:

$$a_\mu^{QED} = 116\,584\,705.7(2.9) \times 10^{-11}, \quad a_\mu^{EW} = 151(4) \times 10^{-11}. \quad (13)$$

4.2 Higher order hadronic contributions to a_μ

Higher order hadronic contributions have been discussed first in [22]. They can be divided into two classes ⁶. The first one involves the vacuum polarization and can be related to the measured $e^+e^- \rightarrow \text{hadrons}$ total cross-section, similar to the lowest order contribution. After rescaling the result in [22, 24], one obtains [1]:

$$a_\mu^{had}(h.o.)_{V.P} = -101.2(6.1) \times 10^{-11}, \quad (14)$$

⁴For more details, see [1].

⁵References to original works can be found in [1, 6, 10].

⁶For more details of the following discussions, see [23].

The second class is the light-by-light scattering diagram. Contrary to the case of vacuum polarizations, this contribution is not yet fully related to a direct measurable quantity. In order to estimate this contribution, one has to introduce some theoretical models. The ones used at present are based on chiral perturbation [25] and ENJL model [26]. To both are added vector meson dominance and phenomenological parametrization of the pion form factors. The different contributions are summarized in Table 4, where the first two come from the boson and quark (constituent) loops, while the last one is due to meson pole exchanges. The first two contributions are quite sensitive to the effects of rho-meson attached at the three off-shell photon legs which reduce the contributions by about one order of magnitude (!). The third one with pseudoscalar meson exchanges (anomaly) gives so far the most important contribution. There is a complete agreement between the two model estimates (after correcting the sign of the pseudoscalar and axial-vector contributions [27]), which may indirectly indicate that the results obtained are model-independent⁷. Nevertheless, there are still some remaining subtle issues to be understood (is the inclusion of a quark loop a double counting ?, why the inclusion of the rho-meson decreases drastically the quark and pion loop contributions ? is a single meson dominance justified?..). The results in Refs. [25] and [26], after correcting the sign of the pseudoscalar and axial-vector contributions as suggested in [27], are given in the table:

Table 4: $a_\mu^{\text{had}}(\text{h.o})_{LL} \times 10^{11}$

Type of diagrams	Ref. [25]	Ref. [26]
π^- loop	-4.5(8.1)	-19(13)
quark loop	9.7(11)	21(3)
π^0, η, η' poles	82.7(6.4)	85(13)
axial-vector pole	1.74	2.5(1.0)
scalar pole	*)	-6.8(2.0)
Total	82.8(15.2)	82.7(18.8)

*) We have added here the result from [26].

while a naive constituent quark model gives [9] using the result of [29]:

$$a_\mu^{\text{had}}(h.o)_{LL} = 92(20) \times 10^{-11} . \quad (15)$$

Due to the unknown real value of the virtual photon momenta entering into the calculation, it can happen that none of the previous approaches describes accurately the LL effect⁸. Therefore, for a conservative estimate, we take an arithmetical average of the three determinations:

$$a_\mu^{\text{had}}(h.o)_{LL} = 84.7(18.0) \times 10^{-11} . \quad (16)$$

One can notice, for the muon, a strong cancellation between the higher order vacuum polarisation and the light by light scattering contributions.

4.3 The total theoretical contributions

Summing up different contributions, the present theoretical status in the standard model is [1]:

$$\begin{aligned} a_\mu^{SM} &= 116\,584\,840.2(19.6) \times 10^{-11} + a_\mu^{\text{had}}(l.o) \\ &= 116\,591\,860.8(78.1) \times 10^{-11} , \end{aligned} \quad (17)$$

⁷See however [28].

⁸I thank Eduardo de Rafael and Paco Yndurain for some clarifying communications on this point.

where $a_\mu^{had}(l.o)$ is the lowest order hadronic contributions evaluated in SN1 [1] (see Eq. (10)). Comparing this SM prediction with the measured value [10]:

$$a_\mu^{exp} = 116\,592\,023(151) \times 10^{-11} , \quad (18)$$

we deduce:

$$a_\mu^{new} \equiv a_\mu^{exp} - a_\mu^{SM} = 162(170) \times 10^{-11} . \quad (19)$$

If the future data will be accurate by $\pm 40 \times 10^{-11}$, while the theoretical errors are almost unchanged due to different limitations discussed previously, then the error in a_μ^{new} will be reduced by a factor 2, giving a chance to detect a 2σ deviation from SM if any. Combined with the mean value of existing determinations of $a_\mu(l.o)^{had}$ given in SN1 [1], which gives: $a_\mu^{new} = 175(170) \times 10^{-11}$, one can deduce the conservative range:

$$-56 \leq a_\mu^{new} \times 10^{11} \leq 393 \quad (90\% \text{ CL}) . \quad (20)$$

4.4 Bounds on some new physics from a_μ

This result gives, for a supersymmetric model with degenerate sparticle mass, a lower bound of about 113 GeV⁹, while the compositeness and the leptoquark scales are constrained to be above 1 TeV. Bound on the sparticle mass is comparable with present experimental bound from LEP data. The one of the leptoquarks is much larger than the present lower bounds of about (200 ~ 300) GeV from direct search experiments at HERA and Tevatron. We expect that these different bounds will be improved in the near future both from accurate measurements of a_μ and of e^+e^- data necessary for reducing the theoretical errors in the determinations of the hadronic contributions, being the major source of the theoretical uncertainties.

4.5 Tau anomaly

In the same way, the higher order hadronic contributions read [1, 31]

$$a_\mu^{had}(h.o)_{VP} = 7.6(0.2) \times 10^{-8} , \quad a_\mu^{had}(h.o)_{LL} = 23.9(5.1) \times 10^{-8} , \quad (21)$$

which, in the τ case, the two effect add each others. Adding the other contributions, one obtains [1]:

$$a_\tau^{SM} = 117\,759.1(6.7) \times 10^{-8} . \quad (22)$$

This value can be compared with the present (inaccurate) experimental one [32]:

$$a_\tau^{exp} = 0.004 \pm 0.027 \pm 0.023 , \quad (23)$$

which, we wish, will be improved in the near future.

5 The QED running coupling $\alpha(M_Z)$

To the lowest order hadronic contribution in Eq. (11), we add the radiative corrections taken by adding the effects of the radiative modes $\pi^0\gamma$, $\eta\gamma$, $\pi^+\pi^-\gamma$, We estimate such effects to be:

$$\Delta\alpha_{had} = (6.4 \pm 2.7) \times 10^{-5} \quad (24)$$

by taking the largest range spanned by the two estimates in [9] and [8]. Using the QED contribution to three-loops [6]:

$$\Delta\alpha_{QED} = 3149.7687 \times 10^{-5} , \quad (25)$$

and the Renormalization Group Evolution of the QED coupling:

$$\alpha^{-1}(M_Z) = \alpha^{-1}(0) \left[1 - \Delta\alpha_{QED} - \Delta\alpha_{had} \right] , \quad (26)$$

one obtains the final estimate:

$$\alpha^{-1}(M_Z) = 128.926(25) , \quad (27)$$

which we show in Fig 1 for a comparison with recent existing determinations. One can notice an improved accuracy of the different recent determinations [6, 8, 9, 20]. We expect that with this new improved estimate of $\alpha(M_Z)$, present lower bound of 114 GeV from LEP data on the Higgs mass can be improved.

⁹For more recent detailed discussions, see e.g. [3, 30] and references quoted there.

6 Muonium hyperfine splitting

Adding to this result in Eq. (12) from SN2 [2], the QED contribution up to fourth order, the lowest order electroweak contribution [7], and an estimate of the higher order weak and hadronic contributions [33]:

$$\begin{aligned} \Delta\nu_{\text{QED}} &= 4\,270\,819(220) \text{ Hz} , \Delta\nu_{\text{weak}}(l.o) = -\frac{G_F}{\sqrt{2}}m_em_\mu\left(\frac{3}{4\pi\alpha}\right)\nu_F \simeq -65 \text{ Hz} , \\ |\Delta\nu_{\text{weak}}(h.o)| &\approx 0.7 \text{ Hz} , \quad \Delta\nu_{\text{had}}(h.o) \simeq 7(2) \text{ Hz} , \end{aligned} \quad (28)$$

one obtains the Standard Model (SM) prediction:

$$\nu_{\text{SM}} \equiv \nu_F + \Delta\nu_{\text{QED}} + \Delta\nu_{\text{weak}} + \Delta\nu_{\text{had}} + \Delta\nu_{\text{had}}(h.o) . \quad (29)$$

If one uses the relation:

$$\nu_F = \rho_F \left(\frac{\mu_\mu}{\mu_B^e} \right) \frac{1}{(1 + m_e/m_\mu)^3} : \quad \rho_F = \frac{16}{3}(Z\alpha)^2 Z^2 cR_\infty , \quad (30)$$

and $Z = 1$ for muonium, $\alpha^{-1}(0) = 137.035\,999\,58(52)$ [10], $cR_\infty = 3\,289\,841\,960\,368(25) \text{ kHz}$ [34], one would obtain:

$$\nu_{\text{SM}} = 4\,463\,302\,913(511)(34)220 \text{ Hz} , \quad (31)$$

where the two first errors are due to the one of the Fermi splitting energy. The first largest one being induced by the one of the ratio of the magnetic moments. The third error is due to the 4th order QED contribution where, one should notice that, unlike the case of a_μ , the dominant errors come from the QED calculation which should then be improved. This prediction can be compared with the precise data [35]:

$$\nu_{\text{exp}} = 4\,463\,302\,776(51) \text{ Hz} . \quad (32)$$

Therefore, at present, we find, it is more informative to extract the Fermi splitting energy ν_F from a comparison of the Standard Model (SM) prediction with the experimental value of ν . Noting that ν_F enters as an overall factor in the theoretical contributions, one can rescale the previous values and predict the ratio:

$$\frac{\nu_{\text{SM}}}{\nu_F} = 1.000\,957\,83(5) . \quad (33)$$

Combining this result with the previous experimental value of ν , one can deduce the SM prediction:

$$\nu_F^{\text{SM}} = 4\,459\,031\,783(226) \text{ Hz} , \quad (34)$$

where the error is dominated here by the QED contribution at fourth order. However, this result is a factor two more precise than the determination in [7] given in Eq. (8), where the main error in Eq. (8) comes from the input values of the magnetic moment ratios. Using this result in Eq. (34) into the expression:

$$\nu_F = \rho_F \left(\frac{m_e}{m_\mu} \right) \frac{1}{(1 + m_e/m_\mu)^3} (1 + a_\mu) , \quad (35)$$

where ρ_F is defined in Eq. (30), and $a_\mu = 1.165\,920\,3(15) \times 10^{-3}$ [10], one can extract a value of the ratio of the muon over the electron mass:

$$\frac{m_\mu}{m_e} = 206.768\,276(11) , \quad (36)$$

to be compared with the PDG value $206.768\,266(13)$ using the masses in MeV units, and with the one from [7]: $206.768\,276(24)$. In Ref. [35], an accuracy two times better than the present result has been also obtained. However, in that case, the errors in the QED contributions may have been underestimated. After inserting the previous value of m_e/m_μ into the alternative (equivalent) relation given in Eq. (30), one can deduce the ratio of magnetic moments:

$$\frac{\mu_\mu}{\mu_B^e} = 4.841\,970\,47(25) \times 10^{-3} , \quad (37)$$

compared to the one obtained from the PDG values of μ_μ/μ_p and μ_p/μ_B^e [14]: $\mu_\mu/\mu_B^e = 4.841\,970\,87(14) \times 10^{-3}$. In both applications, the results in Eqs. (36) and (37) are in excellent agreement with the PDG values. These remarkable agreements can give strong constraints to some contributions beyond the Standard Model and are interesting to be explored.

7 Conclusions

We have evaluated the lowest order hadronic and QCD contributions $a_l^{had}(l.o)$, $\Delta\alpha_{had}$ and $\Delta\nu_{had}$ respectively to the anomalous magnetic moment, QED running coupling and to the muonium hyperfine splitting. Our self-contained results given in Eqs. (10), (11) and (12), obtained within the same strategy and data input, are in excellent agreement with existing determinations and are quite accurate. We have also revised the estimate of the light by light scattering contributions to $a_{\mu,\tau}$, and have explored some phenomenological consequences of these results.

References

- [1] (SN1) S. Narison, *Phys. Lett. B* **513**, 53 (2001); Erratum-ibid. B **256**, 414 (2002) [hep-ph/0103199, version 5] and references therein.
- [2] (SN2) S. Narison, hep-ph/0108065.
- [3] John Ellis, these proceedings and references therein.
- [4] Elena Perazzi, these proceedings.
- [5] M. Gourdin and E. de Rafael, *Nucl. Phys. B* **10**, 667 (1969).
- [6] (J 01) F. Jegerlehner, hep-ph/0104304 and references therein; W.J. Marciano and B.L. Roberts, hep-ph/0105056.
- [7] (CEK 01) A.Czarnecki, S. Eidelman and S.G. Karshenbiom, hep-ph/0107327 and references therein.
- [8] (ADH 98) R. Alemany, M. Davier and A. Höcker, *Eur. Phys. J. C* **2**, 123 (1998).
- [9] (YT 01) J.F. de Trocóniz and F.J. Yndurain, hep-ph/0111258; hep-ph/0107318; hep-ph/0107318.
- [10] The muon g-2 collaboration, H.N. Brown et al., *Phys. Rev. Lett.* **86**, 2227 (2001); A.Czarnecki and W.J. Marciano *Phys. Rev. D* **86**, 013014 (2001); V.W. Hughes and T. Kinoshita, *Rev. Mod. Phys.* **71**, **2**, S133 (1999).
- [11] J.E. Palomar, hep-ph/0202203.
- [12] The ALEPH collaboration, R. Barate et al., *Eur. Phys. J. C* **76**, 15 (1997); C **4**, 409 (1998); A. Hocker, hep-ex/9703004.
- [13] The OPAL collaboration, K. Ackerstaff et al., *Eur. Phys. J. C* **7**, 571 (1999).
- [14] PDG 2000, D.E. Groom et al., *Eur. Phys. J. C* **15**, 1 (2000).
- [15] S. Dolinsky et al., *Phys. Rep. C* **202**, 99 (1991).
- [16] The DM2 collaboration, A. Antonelli et al., *Z. Phys. C* **56**, 15 (1992); D. Bisello et al., *Z. Phys. C* **39**, 13 (1988).
- [17] The DM1 collaboration, F. Mane et al., *Phys. Lett. B* **112**, 178 (1982); A. Cordier et al., *Phys. Lett. B* **110**, 335 (1982).
- [18] The CMD collaboration, R.R. Akhmetshin et al., *Nucl. Phys. A* **675**, 424c (2000); S.I. Serednyakov, *Nucl. Phys. (Proc. Suppl.) B* **96**, 197 (2001).
- [19] The BES collaboration, J.Z. Bai et al., *Phys. Rev. Lett.* **84**, 594 (2000); hep-ex/0102003.
- [20] (DH 98) M. Davier and A. Hoecker, *Phys. Lett. B* **435**, 419 (1998); (MOR 01) A.D. Martin, J. Outhwaite and M.G. Ryskin, hep-ph/0012231; (BP 01) H. Burkhardt and B. Pietrzyk, LAPP-EXP 2001-03 (2001).
- [21] (FKM 99) R.N. Faustov, A. Karimkhodzhaev and A.P. Martynenko, *Phys. Rev. A* **59**, 2498 (1999).
- [22] J. Calmet, S. Narison, M. Perrottet and E. de Rafael, *Rev. Mod. Phys.* **49**, 21 (1977); *Phys. Lett. B* **161**, 283 (1976); S. Narison, *Thèse de 3ème cycle*, Marseille (1976).
- [23] S. Narison, *QCD as a theory of hadrons*, Cambridge Univ. Press (to appear).
- [24] B. Krause, *Phys. Lett. B* **390**, 392 (1997).
- [25] M. Hayakawa and T. Kinoshita, *Phys. Rev. D* **57**, 465 (1998); hep-ph/0112102.
- [26] J. Bijnens, E. Pallante and J. Prades, *Nucl. Phys. B* **474**, 379 (1996); hep-ph/0112255.

- [27] M. Knecht and A. Nyffeler, hep-ph/0111058; M. Knecht, A. Nyffeler, M. Perrottet and E. de Rafael, hep-ph/0111059; see also: [25, 26] and I. Blokland, A. Czarnecki and K. Melnikov, hep-ph/0112117.
- [28] M. Ramsey-Musolf and M.B. Wise, hep-ph/0112255.
- [29] S. Laporta and E. Remiddi, *Phys. Lett. B* **301**, 440 (1993).
- [30] U. Chattopadhyay, A. Corsetti and P. Nath, hep-ph/0202275.
- [31] S. Narison, *J. Phys. (Nucl. Phys.) G* **4**, 1840 (1978).
- [32] L. Taylor, *Nucl. Phys. (Proc. Suppl.) B* **76**, 273 (1999).
- [33] S.G. Karshenboim, *Z. Phys. D* **36**, 11 (1996).
- [34] P.J. Mohr and B.N. Taylor, *Rev. Mod. Phys.* **72**, 351 (2000).
- [35] W. Liu et al., *Phys. Rev. Lett.* **82**, 711 (1999); F.G. Mariam et al., *Phys. Rev. Lett.* **49**, 993 (1982).